Stability of self-gravitating accretion disks in galactic centers Lumpy gravy

Markward Britsch

Supervisor: Wolfgang J. Duschl

Markward.Britsch@ita.uni-heidelberg.de

Zentrum für Astronomie Heidelberg, Institut für Theoretische Astrophysik Universität Heidelberg







How to fix lumpy gravy

No matter how many times you make pan gravy, lumps have a way of appearing—usually when you least want them.

http://www.ehow.com/how_114366_fix-lumpy-gravy.html







How to fix lumpy gravy

No matter how many times you make pan gravy, lumps have a way of appearing—usually when you least want them.

http://www.ehow.com/how_114366_fix-lumpy-gravy.html

At least on larger galactic scales, it is believed that strong self-gravity leads to star formation. If the same is true of QSO discs, there is a danger that most of the gas would form stars, leaving little to fuel the QSO.

Goodman (2003)







How to fix lumpy gravy

No matter how many times you make pan gravy, lumps have a way of appearing—usually when you least want them.

Don't throw the gravy out. [...] You can still put Thanksgiving dinner on the table in time.

http://www.ehow.com/how_114366_fix-lumpy-gravy.html

At least on larger galactic scales, it is believed that strong self-gravity leads to star formation. If the same is true of QSO discs, there is a danger that most of the gas would form stars, leaving little to fuel the QSO.

Goodman (2003)








A SMBH formation model

merger of two galaxies → gas flows into the center












A SMBH formation model

- merger of two galaxies → gas flows into the center
- it forms an accretion disk that feeds the forming black hole











• such a disk is gravitational unstable if it is not hot








- such a disk is gravitational unstable if it is not hot
- simulate such a thin disk in polar coordinates















- such a disk is gravitational unstable if it is not hot
- simulate such a thin disk in polar coordinates
- $R_{\rm i} = 29 {
 m pc}, R_{\rm a} = 126 {
 m pc},$ $M_{\rm d} = 6 \cdot 10^8 M_{\odot} \gg M_* = 7.5 \cdot 10^7 M_{\odot}$
- some viscosity, dynamically not important since $t_{\nu} \gg t_{\rm end} = 3.2 \cdot 10^6 {\rm a}$







- such a disk is gravitational unstable if it is not hot
- simulate such a thin disk in polar coordinates
- $R_{\rm i} = 29 {
 m pc}, R_{\rm a} = 126 {
 m pc},$ $M_{\rm d} = 6 \cdot 10^8 M_{\odot} \gg M_* = 7.5 \cdot 10^7 M_{\odot}$
- some viscosity, dynamically not important since $t_{\nu} \gg t_{\rm end} = 3.2 \cdot 10^6 {\rm a}$
- use NIRVANA2.0 (Ziegler, 1998, 1999)







- such a disk is gravitational unstable if it is not hot
- simulate such a thin disk in polar coordinates
- $R_{\rm i} = 29 {
 m pc}, R_{\rm a} = 126 {
 m pc},$ $M_{\rm d} = 6 \cdot 10^8 M_{\odot} \gg M_* = 7.5 \cdot 10^7 M_{\odot}$
- some viscosity, dynamically not important since $t_{\nu} \gg t_{\rm end} = 3.2 \cdot 10^6 {\rm a}$
- use NIRVANA2.0 (Ziegler, 1998, 1999)
- include cooling







- such a disk is gravitational unstable if it is not hot
- simulate such a thin disk in polar coordinates
- $R_{\rm i} = 29 {
 m pc}, R_{\rm a} = 126 {
 m pc},$ $M_{\rm d} = 6 \cdot 10^8 M_{\odot} \gg M_* = 7.5 \cdot 10^7 M_{\odot}$
- some viscosity, dynamically not important since $t_{\nu} \gg t_{\rm end} = 3.2 \cdot 10^6 {\rm a}$
- use NIRVANA2.0 (Ziegler, 1998, 1999)
- include cooling
- no star formation







- such a disk is gravitational unstable if it is not hot
- simulate such a thin disk in polar coordinates
- $R_{\rm i} = 29 {
 m pc}, R_{\rm a} = 126 {
 m pc},$ $M_{\rm d} = 6 \cdot 10^8 M_{\odot} \gg M_* = 7.5 \cdot 10^7 M_{\odot}$
- some viscosity, dynamically not important since $t_{\nu} \gg t_{\rm end} = 3.2 \cdot 10^6 {\rm a}$
- use NIRVANA2.0 (Ziegler, 1998, 1999)
- include cooling
- no star formation










Accreted mass









• more simulations done with different disk masses and values for the viscosity







- more simulations done with different disk masses and values for the viscosity
- high accretion due to clump-clump interactions $t_{\rm acc} \propto \frac{1}{\Omega}$







- more simulations done with different disk masses and values for the viscosity
- high accretion due to clump-clump interactions $t_{\rm acc} \propto \frac{1}{\Omega}$
- as predicted by the β -viscosity (Duschl, Strittmatter, Biermann, 2000 and Duschl, Britsch submitted to ApJL)







- more simulations done with different disk masses and values for the viscosity
- high accretion due to clump-clump interactions $t_{\rm acc} \propto \frac{1}{\Omega}$
- as predicted by the β -viscosity (Duschl, Strittmatter, Biermann, 2000 and Duschl, Britsch submitted to ApJL)
- turbulent velocities and length-scales also match the β -viscosity







- more simulations done with different disk masses and values for the viscosity
- high accretion due to clump-clump interactions $t_{\rm acc} \propto \frac{1}{\Omega}$
- as predicted by the β -viscosity (Duschl, Strittmatter, Biermann, 2000 and Duschl, Britsch submitted to ApJL)
- turbulent velocities and length-scales also match the β -viscosity
- $t_{\rm acc} \propto \frac{1}{\Omega}$ also predicted by a clumpy disk model by Kumar (1999)







Summary

- formation model of SMBHs including large thin self-gravitating disks
- interacting clumps give very high accretion rate
- can be interpreted as a β -viscosity
- exciting accretion process
- supports this SMBH formation model
- but there's still a lot to do ...







References

Goodman, J. MNRAS 339 (2003) 937-948

Duschl, W. J., Strittmatter, P. A., Biermann, P. L A&A

357 (2000) 1123-1132

Duschl, W. J., Strittmatter, P. A.,

arXiv:astro-ph/0602009 (2006)

Barnes, J. E., Hernquist, L., ApJ 471 (1996) 115-142

Ziegler, U., Comp. Phys. Comm. 109 (1998) 111-134

Ziegler, U., Comp. Phys. Comm. 116 (1999) 65-77

Kumar, P., ApJ 519 (1999) 599-604





